

*The following resources related to this article are available online at [jada.ada.org](http://jada.ada.org) ( this information is current as of November 20, 2009 ):*

**Updated information and services** including high-resolution figures, can be found in the online version of this article at:

<http://jada.ada.org/cgi/content/full/131/11/1559>

This article appears in the following **subject collections**:

Imaging <http://jada.ada.org/cgi/collection/imaging>

Information about obtaining **reprints** of this article or about permission to reproduce this article in whole or in part can be found at:

<http://www.ada.org/prof/resources/pubs/jada/permissions.asp>

# COVER STORY

ROBERT A. FREITAS JR., J.D., B.S.

## ABSTRACT

**Background.** Nanodentistry will make possible the maintenance of comprehensive oral health by involving the use of nanomaterials, biotechnology (including tissue engineering) and, ultimately, dental nanorobotics (nanomedicine).

**Results.** When the first micrometer-sized dental nanorobots can be constructed within 10 to 20 years, these devices will allow precisely controlled oral analgesia, dentition replacement therapy using biologically autologous whole replacement teeth man-

ufactured during a single office visit, and rapid nanometer-scale precision restorative dentistry.

**Clinical Implications.** New treatment opportunities may include dentition renaturalization, permanent hypersensitivity cure, complete orthodontic realignments during a single office visit, covalently bonded diamondized enamel and continuous oral health maintenance through the use of mechanical dentifrobots.

# NANODENTISTRY

In 1959, the late Nobel Prize-winning physicist Richard P. Feynman presented a talk entitled "There's Plenty of Room at the Bottom"<sup>1</sup> at the annual meeting of the American Physical Society. Feynman proposed using machine tools to make smaller machine tools, which, in turn, would be used to make still smaller machine tools, and so on all the way down to the molecular level. He suggested that such nanomachines, nanorobots and nanodevices ultimately could be used to develop a wide range of atomically precise microscopic instrumentation and manufacturing tools. Feynman<sup>1</sup> argued that these tools could be applied to produce vast quantities of ultrasmall computers and various microscale and nanoscale robots. He concluded that this is "a development which I think cannot be avoided." The vision of nanotechnology was born.

Forty years ago, this talk was greeted with astonishment and skepticism. However, since then, we have made remarkable progress toward realizing Feynman's vision. From the dawn of the microcomputer era several decades ago, we have witnessed a significant increase in the speed and power of computers. This is due, in large measure, to the ever-decreasing size of the electronic components that can be packed at ever-increasing densities onto a single silicon chip. Transistor density has doubled every 18 months, an observation that has come to be

known as Moore's law.<sup>2</sup> The size of features on computer chips has shrunk from a fraction of a millimeter in the first microprocessor chip to 0.1 to 0.2 micrometers ( $1 \mu\text{m} = 10^{-6}$  meter) in the latest chips. Recently, two new companies have been formed<sup>3,4</sup> with the explicit goal of producing molecular computer components<sup>4,5</sup> using molecular parts at the nanometer ( $10^{-9}$  meter, or one-billionth of a meter) scale, within just a few years.

Similar progress is under way in the related field of robotic miniaturization. The burgeoning field of microelectromechanical systems was made possible by the fabrication of the first micromotors in the late 1980s and early 1990s.<sup>6</sup> By 1994, engineers at Nippondenso Ltd.<sup>7</sup> had constructed a working electric car smaller than a grain of rice, a 1/1,000th-scale replica of a 1936 Model AA Toyota sedan that incorporated 24 parts, including a motor, wheels, body, spare tire, bumpers and even a 10- $\mu\text{m}$ -thick license plate. In 1997, researchers at Cornell's Nanofabrication Facility<sup>8</sup> produced a silicon guitar that was 10  $\mu\text{m}$  in length and 2  $\mu\text{m}$  wide, with six individual "strings" that were only 50 nanometers (approximately 200 atoms) thick.

Even Feynman had notions of how nanotechnology could be applied to medicine. After discussing his ideas with a colleague, Feynman<sup>1</sup> offered the first known proposal for a nanomedical procedure to cure heart disease: "A friend of

mine [Albert R. Hibbs] suggests an interesting possibility for relatively small machines. He says that, although it is a very wild idea, it would be interesting in surgery if you could swallow the surgeon. You put the mechanical surgeon inside the blood vessel and it goes into the heart and looks around. (Of course, the information has to be fed out.) It finds out which valve is the faulty one and takes a little knife and slices it out. Other small machines might be permanently incorporated in the body to assist some inadequately functioning organ.”


Once one considers other potential applications of nanotechnology to medicine, it is not difficult to imagine what nanodentistry would look like. The main purpose of this report, therefore, is to provide an early glimpse of nanodental applications and to illustrate their potentially far-reaching impact on clinical dental practice. In this article, I briefly survey the field of nanomedicine and present some potential applications to dentistry, such as local anesthesia, reconstruction of dental hard tissues, orthodontic treatment and disease prevention. I conclude by placing the expected development of nanodentistry, which may see its earliest practical uses within the next 10 to 20 years, in the context of today's trends in dental science and practice.

#### NANOMEDICINE

Molecular manufacturing<sup>9,10</sup> promises precise control of matter at the atomic and molecular level. One major implication of this is that, within the next 10 to 20 years, it should become possible to construct machines

on the micrometer scale made up of parts on the nanometer scale. Subassemblies of such devices may include such useful robotic components as 100-nm manipulator arms, 10-nm sort-

Other applications include mechanically reversing atherosclerosis,<sup>17</sup> improving respiratory capacity,<sup>18</sup> enabling near-instantaneous hemostasis,<sup>19</sup> supplementing the immune sys-



ing rotors for molecule-by-molecule reagent purification, and smooth superhard surfaces made of atomically flawless diamond (a durable, well-characterized material, composed of abundant carbon atoms, that should be relatively easy to manufacture).<sup>10</sup>

Nanocomputers would assume the important task of activating, controlling and deactivating such nanomechanical devices. Nanocomputers would store and execute mission plans, receive and process external signals and stimuli, communicate with other nanocomputers or external control and monitoring devices, and possess contextual knowledge to ensure safe functioning of the nanomechanical devices.

Such technology has enormous medical implications.<sup>11</sup> Programmable nanorobotic devices would allow physicians to perform precise interventions at the cellular and molecular level. Medical nanorobots have been proposed for gerontological applications,<sup>12</sup> in pharmaceutical research<sup>13</sup> and clinical diagnosis,<sup>11,14</sup> and in dentistry.<sup>15,16</sup>

tem,<sup>15,20</sup> rewriting<sup>21</sup> or replacing<sup>22</sup> DNA sequences in cells, repairing brain damage<sup>23</sup> and resolving gross cellular insults,<sup>21</sup> whether caused by “irreversible” processes<sup>24</sup> or by cryogenic storage of biological tissues.<sup>9,25</sup>

Growing interest in the future medical applications of nanotechnology is leading to the emergence of a new field called nanomedicine.<sup>11</sup> This is the science and technology of diagnosing, treating and preventing disease and traumatic injury; of relieving pain; and of preserving and improving human health, through the use of nanoscale-structured materials, biotechnology and genetic engineering, and eventually complex molecular machine systems and nanorobots. Similarly, development of nanodentistry will make possible the maintenance of near-perfect oral health through the use of nanomaterials<sup>26,27</sup>; biotechnology,<sup>27-31</sup> including tissue engineering<sup>32,33</sup>; and nanorobotics. The first two of these approaches have been discussed extensively elsewhere. This report offers a first

look at the third—and most exciting (although somewhat more technologically remote)—approach to nanodentistry: dental nanorobotics.

#### APPLICATIONS OF NANOROBOTICS TO DENTISTRY

When the first micrometer-sized dental nanorobots are constructed, perhaps 10 to 20 years from today, how might they be applied to dentistry? I<sup>11</sup> have described how medical nanorobots might use specific motility mechanisms to crawl or swim through human tissues with navigational precision; acquire energy, and sense and manipulate their surroundings; achieve safe cytopenetration (for example, pass through plasma membranes such as the odontoblastic process without disrupting the cell, while maintaining clinical biocompatibility<sup>11</sup>); and use any of a multitude of techniques to monitor, interrupt or alter nerve-impulse traffic in individual nerve cells.

These nanorobotic functions may be controlled by an onboard nanocomputer that executes preprogrammed instructions in response to local

robots via acoustic signals (as are used in ultrasonography) or other means—similar to an admiral commanding a fleet.


**Inducing anesthesia.** One of the most common procedures in dentistry is the injection of local anesthetic, which can involve long waits and varying degrees of efficacy, patient discomfort and complications.<sup>34</sup> Well-known alternatives, such as transcutaneous electronic nerve stimulation,<sup>35,36</sup> cell demodulated electronic targeted anesthesia<sup>37</sup> and other transmucosal,<sup>36</sup> intraosseous<sup>36</sup> or topical<sup>38</sup> techniques, are of limited clinical effectiveness.

To induce oral anesthesia in the era of nanodentistry, dental professionals will instill a colloidal suspension containing millions of active analgesic micrometer-sized dental nanorobot “particles” on the patient’s gingivae. After contacting the surface of the crown or mucosa, the ambulating nanorobots reach the dentin by migrating into the gingival sulcus and passing painlessly through the lamina propria<sup>39</sup> or the 1- to 3- $\mu\text{m}$ -thick layer of loose tissue at the cementodentinal junction.<sup>40</sup> On reaching

guided by a combination of chemical gradients, temperature differentials and even positional navigation,<sup>11</sup> all under the control of the onboard nanocomputer, as directed by the dentist.

There are many pathways to choose from. Dentinal tubule number density is typically 22,000  $\text{mm}^{-2}$  near the dentino-enamel junction, 37,000  $\text{mm}^{-2}$  midway between the junction and the pulpal wall, and 48,000  $\text{mm}^{-2}$  close to the pulp<sup>41</sup> in coronal dentin,<sup>41</sup> with the number density slightly lower in the root (for example, 13,000  $\text{mm}^{-2}$  near the cementum<sup>44</sup>). Tubule diameter increases nearer the pulp,<sup>43,45</sup> which may facilitate nanorobot movement, although circumpulpal tubule openings vary in number and size.<sup>46</sup>

Tubule branching patterns may present a significant challenge to navigation, because they exhibit an intricate and profuse canalicular anastomosing system that crisscrosses the intertubular dentin, with dentinal branching density most abundant in locations where tubule density is low.<sup>47</sup> Dentinal tubules are continuous between primary dentin and regular secondary dentin in young and old teeth, but not between primary and irregular secondary dentin.<sup>48</sup> Regular secondary dentin becomes highly sclerosed in older teeth,<sup>48</sup> and many tubule openings on the outer dentin surface can become completely occluded in some circumstances,<sup>49</sup> probably requiring significant detouring by the dental nanorobots. (On the other hand, a small number of microcanals,<sup>50,51</sup> large tubules<sup>52</sup> or giant tubules<sup>51,53</sup> with diameters of 10 to 50  $\mu\text{m}$  or even larger<sup>54</sup> may exist in some




**Tubule branching patterns may present a significant challenge to navigation of nanorobots.**

sensor stimuli. Alternatively, the dentist may issue strategic instructions by transmitting orders directly to in vivo nano-

the dentin, the nanorobots enter dentinal tubule holes that are 1 to 4  $\mu\text{m}$  in diameter<sup>41-43</sup> and proceed toward the pulp,

cases, possibly affording easier transit.)

Assuming a total path length of about 10 mm from the tooth surface to the pulp and a modest travel speed<sup>11</sup> of 100  $\mu\text{m/s}$ ,



nanorobots can complete the journey into the pulp chamber in approximately 100 seconds. The presence of natural cells that are constantly in motion around and inside the teeth—including human gingival and pulpal fibroblasts,<sup>55</sup> cementoblasts at the cementodentinal junction,<sup>56</sup> bacteria inside dentinal tubules,<sup>57</sup> odontoblasts near the pulpal/dentin border<sup>43,46,58,59</sup> and lymphocytes within the pulp or lamina propria<sup>60</sup>—suggests that such journeys should be feasible by cell-sized nanorobots of similar mobility.

Once installed in the pulp and having established control over nerve-impulse traffic,<sup>11</sup> the analgesic dental nanorobots may be commanded by the dentist to shut down all sensitivity in any tooth that requires treatment. When the dentist presses the icon for the desired tooth on the hand-held controller display, the selected tooth immediately numbs (or later, on command, awakens). After the oral procedures are completed, the dentist orders the nanorobots (via the same acoustic data links) to restore all sensation, to relinquish control of nerve

traffic and to egress from the tooth via similar pathways used for ingress; following this, they are aspirated. Nanorobotic analgesics offer greater patient comfort and reduced anxiety,

no needles,<sup>61,62</sup> greater selectivity and controllability of the analgesic effect, fast and completely reversible action, and avoidance of most side effects and complications.<sup>34</sup>

New treatment opportunities abound in nanodentistry, as illustrated by these examples:

#### **Major tooth repair.**

Nanodental techniques for major tooth repair may evolve through several stages of technological development, first using genetic engineering, tissue engineering<sup>32,33,63-67</sup> and tissue regeneration,<sup>33,63-68</sup> and later involving the growth of whole new teeth in vitro<sup>69,70</sup> and their installation. Ultimately, the nanorobotic manufacture and installation of a biologically autologous whole-replacement tooth that includes both mineral and cellular<sup>71-73</sup> components—that is, complete dentition replacement therapy—should become feasible within the time and economic constraints of a typical office visit, through the use of an affordable desktop manufacturing facility, which would fabricate the new tooth, in the dentist's office.

#### **Renaturalization pro-**

**cedures.** Dentition renaturalization procedures may become a popular addition to the typical dental practice, providing perfect treatment methods for esthetic dentistry. This trend may begin with patients who desire to have their old dental amalgams<sup>74</sup> excavated and their teeth remanufactured with native biological materials. However, demand will grow for full coronal renaturalization procedures in which all fillings, crowns and other 20th-century modifications to the visible dentition are removed, with the affected teeth remanufactured to become indistinguishable from the original teeth.

#### **Dentin hypersensitivity.**

Dentin hypersensitivity is another pathological phenomenon that may be amenable to nanodental treatment. Dentin hypersensitivity may be caused by changes in pressure transmitted hydrodynamically to the pulp. This etiology is suggested by the finding that hypersensitive teeth have dentinal tubules with surface number densities that are eight times higher than those of nonsensitive teeth, as well as tubules with diameters that are twice as large.<sup>75</sup> Many therapeutic agents provide temporary relief for this common painful condition,<sup>76</sup> but reconstructive dental nanorobots, using native biological materials, could selectively and precisely occlude specific tubules within minutes, offering patients a quick and permanent cure.

#### **Tooth repositioning.**

Orthodontic nanorobots could directly manipulate the periodontal tissues, including gingivae, periodontal ligament, cementum and alveolar bone, allowing rapid and painless

tooth straightening, rotating and vertical repositioning within minutes to hours. This is in contrast to current molar-uprighting techniques, which require weeks or months to complete.<sup>77</sup>

**Durability and appearance.** Tooth durability and appearance may be improved by replacing upper enamel layers with covalently bonded artificial materials such as sapphire<sup>78</sup> or diamond,<sup>16</sup> which have 20 to 100 times the hardness and failure strength (that is, the pressure that must be applied to cause a solid material to fail catastrophically) of natural enamel<sup>11</sup> or contemporary ceramic veneers,<sup>79</sup> as well as good biocompatibility.<sup>80</sup> Like enamel, sapphire is somewhat susceptible to acid corrosion,<sup>81</sup> but sapphire can be manufactured in virtually any color of the rainbow,<sup>11</sup> offering interesting cosmetic alternatives (for example, iridescence) to standard whitening<sup>82</sup> and sealant<sup>83</sup> procedures. Pure sapphire and diamond are brittle and prone to fracture if sufficient shear forces are imposed,<sup>84</sup> but they can be made more fracture-resistant as part of a nanostructured composite material<sup>85</sup> that possibly includes embedded carbon nanotubes.<sup>86</sup>

Effective prevention has reduced the incidence of caries in children<sup>87</sup> and a caries vaccine may soon be available.<sup>88</sup> However, a subocclusal-dwelling nanorobotic dentifrice delivered by mouthwash or toothpaste could patrol all supragingival and subgingival surfaces at least once a day, metabolizing trapped organic matter into harmless and odorless vapors and performing continuous calculus débridement.


These almost-invisible (1 to 10  $\mu\text{m}$ ) dentifrobots, perhaps numbering  $10^3$  to  $10^5$  per mouth and crawling at 1 to 10  $\mu\text{m}/\text{second}$ , might have the mobility of tooth amoebas,<sup>89</sup> but would be inexpensive, purely mechanical devices that safely deactivate themselves if swallowed. Moreover, they would be programmed with strict protocols to avoid occlusal surfaces. (Even diamondoid nanorobots can be crushed by dental grinding unless the thickness of their outer shells is at least 10 percent of the device's radius.<sup>11</sup>)

Properly configured dentifrobots could identify and destroy pathogenic bacteria residing in the plaque and elsewhere, while allowing the 500 or so species of harmless oral microflora to flourish in a healthy ecosystem. Dentifrobots also would provide a continuous barrier to halitosis, since bacterial putrefaction is the central metabolic process involved in oral malodor.<sup>90</sup> With this kind of daily dental care available

theoretical and applied research needed to turn them into reality is progressing rapidly. Nanotechnological developments are expected to accelerate significantly through new governmental<sup>91</sup> and private-sector<sup>92</sup> initiatives.

Nanotechnological advances should be viewed in the context of other expected developments relevant to oral health in the coming decades. Biological approaches such as tissue and genetic engineering<sup>32,33,63-67</sup> will yield new diagnostic and therapeutic approaches much sooner than will nanotechnology. At the same time, continual refinement of traditional methods, development of advanced restorative materials, and new medications and pharmacological approaches will continue to improve dental care.

Trends in oral health and disease also may change the focus on specific diagnostic and treatment modalities. Increasingly preventive approaches will reduce the need for cura-



Properly configured dentifrobots could identify and destroy pathogenic bacteria residing in the plaque and elsewhere.

from an early age, conventional tooth decay and gingival disease will disappear.

#### THE PATH TO NANODENTISTRY

The visions described thus far may sound unlikely, implausible or even heretic. Yet, the

tive or restorative interventions, as has already happened with dental caries. Deeper understanding of the causes and pathogenesis of other disease processes—such as periodontal disease, developmental craniofacial defects and malignant neoplasms—should make

prevention a viable approach for most of them.

The role of the dentist will continue to evolve along the lines of currently visible trends. In the United States, for example, cases involving simple self-care neglect will become fewer, while cases involving cosmetic procedures, acute trauma or rare disease conditions will become relatively more commonplace. Diagnosis and treatment will be customized to match the preferences and genetics of each patient. Treatment options will become more numerous and exacting. All this will demand, even more so than today, the best technical abilities, professional judgment and strong interpersonal skills that are the hallmark of the contemporary dentist.

## CONCLUSION

Nanodentistry still faces many significant challenges<sup>11</sup> in realizing its tremendous potential. Basic engineering problems run the gamut from precise positioning and assembly of molecular-scale parts, to economical mass-production techniques, to biocompatibility and the simultaneous coordination of the activities of large numbers of independent micrometer-scale robots. In addition, there are larger social issues of public acceptance, ethics, regulation and human safety that must be addressed before molecular nanotechnology can enter the modern medical armamentarium. However, there are equally powerful motivations to surmount these various challenges, such as the possibility of providing high-quality dental care to the 80 percent of the world's population that currently receives no significant dental

care. Time, specific advances, financial and scientific resources, and human needs will determine which of the applications described in this article are realized first. ■

Mr. Freitas is a research scientist with Zyvex Corp., a nanotechnology research and development company, 1321 N. Plano Road, Richardson, Texas 75081, e-mail "rfreitas@calweb.com". Address reprint requests to Mr. Freitas.

The author thanks Drs. Titus Schleyer and Heiko Spallek, as well as several anonymous reviewers, for helpful comments on the manuscript.

1. Feynman RP. There's plenty of room at the bottom. *Eng Sci Feb.* 1960;23:22-36. Available at: "www.zyvex.com/nanotech/feynman.html". Accessed Sept. 26, 2000.

2. Schleyer TK. Digital dentistry in the computer age. *JADA* 1999;130:1713-20.

3. California Molecular Electronics Corporation (CALMEC). Available at: "www.calmec.com". Accessed Sept. 26, 2000.

4. Reed MA, Tour JM. Computing with molecules. *Sci Am* 2000;282(6):86-93. Markoff J.

5. Collier CP, Wong EW, Belohradsky M, et al. Electronically configurable molecular-based logic gates. *Science* 1999;285(5426):391-4.

6. Mehregany M, Tai Y-C. Surface micro-machined mechanisms and micromotors. *J Micromech Microeng* 1991;1:73-85.

7. Teshigahara A, Watanabe M, Kawahara N, Ohtsuka Y, Hattori T. Performance of a 7-mm microfabricated car. *J Microelectromech Sys* 1995;4:76-80.

8. Carr DW, Craighead HG. Fabrication of nanoelectromechanical systems in single crystal silicon using silicon on insulator substrates and electron beam lithography. *J Vac Sci Technol B* 1997;15:2760-5.

9. Drexler KE. Molecular engineering: an approach to the development of general capabilities for molecular manipulation. *Proc Natl Acad Sci USA* 1981;78(9):5275-8. Available at: "www.imm.org/PNAS.html". Accessed Sept. 26, 2000.

10. Drexler KE. *Nanosystems: Molecular machinery, manufacturing, and computation*. New York: Wiley; 1992.

11. Freitas RA Jr. *Nanomedicine*. Vol. 1. Basic capabilities. Georgetown, Texas: Landes Bioscience; 1999. Available at: "www.nanomedicine.com". Accessed Sept. 26, 2000.

12. Fahy GM. Short-term and long-term possibilities for interventive gerontology. *Mt Sinai J Med* 1991;58(4):328-40.

13. Fahy GM. Molecular nanotechnology and its possible pharmaceutical implications. In: Bezold C, Halperin JA, Eng JL, eds. *2020 visions: Health care information standards and technologies*. Rockville, Md.: U.S. Pharmacopeial Convention; 1993:152-9.

14. Lampton C. Nanotechnology promises to revolutionize the diagnosis and treatment of diseases. *Genet Eng News* 1995;15:4. 23.

15. Drexler KE, Peterson C, Pergamit G. Unbounding the future: The nanotechnology

revolution. New York: William Morrow/Quill Books; 1991:199-225.

16. Reifman EM. Diamond teeth. In: Crandall BC, ed. *Nanotechnology: Molecular speculations on global abundance*. Cambridge, Mass.: MIT Press; 1996:81-6.

17. Dewdney AK. Nanotechnology: wherein molecular computers control tiny circulatory submarines. *Sci Am* 1988;258(1):100-3.

18. Freitas RA Jr. Exploratory design in medical nanotechnology: a mechanical artificial red cell. *Artif Cells Blood Substit Immobil Biotechnol* 1998;26(4):411-30. Available at: "www foresight.org/Nanomedicine/Respirocytes.html". Accessed Sept. 26, 2000.

19. Freitas RA Jr. Clotocytes: artificial mechanical platelets. *Foresight Update* 2000;41(June 30):9-11. Available at: "www.imm.org/Reports/Rep018.html". Accessed Sept. 26, 2000.

20. Merkle RC. Nanotechnology and medicine. In: Klatz R, Kovarik FA, Goldman B, eds. *Advances in anti-aging medicine*. Vol. 1. Larchmont, N.Y.: Mary Ann Liebert; 1996:277-86. Available at: "www.zyvex.com/nanotech/nanotechAndMedicine.html". Accessed Sept. 26, 2000.

21. Drexler KE. *Engines of creation: The coming era of nanotechnology*. New York: Anchor Press/Doubleday; 1986:99-129. Available at: "www foresight.org/EOC/". Accessed Sept. 26, 2000.

22. Freitas RA Jr. Say Ahhh! The Sciences 2000;40(July/August):26-31. Available at: "www foresight.org/Nanomedicine/SayAh/index.html". Accessed Sept. 26, 2000.

23. Merkle RC. The molecular repair of the brain. Available at: "www.merkle.com/cryo/techFeas.html". Accessed Sept. 26, 2000.

24. Fahy GM. Possible medical applications of nanotechnology. In: Crandall BC, Lewis J, eds. *Nanotechnology: Research and perspectives*. Cambridge, Mass.: MIT Press; 1992:251-67.

25. Merkle RC. The technical feasibility of cryonics. *Med Hypotheses* 1992;39(1):6-16.

26. Shi H, Tsai WB, Garrison MD, Ferrari S, Ratner BD. Template-imprinted nanostructured surfaces for protein recognition. *Nature* 1999;398(6728):593-7.

27. West JL, Halas NJ. Applications of nanotechnology to biotechnology. *Curr Opin Biotechnol* 2000;11(2):215-7.

28. Pruzansky S. Effect of molecular genetics and genetic engineering on the practice of orthodontics. *Am J Orthod* 1972;62(5):539-42.

29. Farr C. Biotech in periodontics: molecular engineering yields new therapies. *Dent Today* 1997;16(10):92, 94-7.

30. Slavkin HC. Entering the era of molecular dentistry. *JADA* 1999;130(3):413-7.

31. Sims MR. Brackets, epitopes and flash memory cards: a futuristic view of clinical orthodontics. *Aust Orthod J* 1999;15(5):260-8.

32. Baum BJ, Mooney DJ. The impact of tissue engineering on dentistry. *JADA* 2000;131:309-18.

33. Buckley MJ, Agarwal S, Gassner R. Tissue engineering and dentistry. *Clin Plast Surg* 1999;26(4):657-62.

34. Justig JP, Zusman SP. Immediate complications of local anesthetic administered to 1,007 consecutive patients. *JADA* 1999;130:496-9.

35. Hochman R. Neurotransmitter modulator (TENS) for control of dental operative pain. *JADA* 1988;116:208-12.

36. Estafan DJ. Invasive and noninvasive

dental analgesia techniques. *Gen Dent* 1998;46(6):600-3.

37. Cedeta Dental International Inc. Clinical Studies, 1991-1997. Available at: "www.cedeta dental.com/clinical.htm". Accessed Sept. 26, 2000.
38. Meechan JG. Intra-oral topical anaesthetics: a review. *J Dent* 2000;28(1):3-14.
39. Paulsen F, Thale A. Epithelial-connective tissue boundary in the oral part of the human soft palate. *J Anat* 1998;193:457-67.
40. Yamamoto T, Doman T, Takahashi S, Islam N, Suzuki R, Wakita M. The structure and function of the cemento-dentinal junction in human teeth. *J Periodontol Res* 1999;34(5):261-8.
41. Dourda AO, Moule AJ, Young WG. A morphometric analysis of the cross-sectional area of dentine occupied by dentinal tubules in human third molar teeth. *Int Endod J* 1994;27(4):184-9.
42. Arends J, Stokroos I, Jongbloed WG, Ruben J. The diameter of dentinal tubules in human coronal dentine after demineralization and air drying: a combined light microscopy and SEM study. *Caries Res* 1995;29(2):118-21.
43. Goracci G, Mori G. Micromorphological aspects of dentin (in Italian). *Minerva Stomatol* 1995;44(9):377-87.
44. Tidmarsh BG, Arrowsmith MG. Dentinal tubules at the root ends of apicect teeth: a scanning electron microscopic study. *Int Endod J* 1989;22(4):184-9.
45. Marchetti C, Piacentini C, Menghini P. Morphometric computerized analysis on the dentinal tubules and the collagen fibers in the dentine of human permanent teeth. *Bull Group Int Rech Sci Stomatol Odontol* 1992;35(3-4):125-9.
46. Marion D, Jean A, Hamel H, Kerebel LM, Kerebel B. Scanning electron microscopic study of odontoblasts and circum-pulpal dentin in a human tooth. *Oral Surg Oral Med Oral Pathol* 1991;72(4):473-8.
47. Mjör IA, Nordahl I. The density and branching of dentinal tubules in human teeth. *Arch Oral Biol* 1996;41(5):401-12.
48. Wang YN, Ashrafi SH, Weber DF. Scanning electron microscopic observations of casts of human dentinal tubules along the interface between primary and secondary dentine. *Anat Rec* 1985;211(2):149-55.
49. Brannstrom M, Garberoglio R. Occlusion of dentinal tubules under superficial attrited dentine. *Swed Dent J* 1980;4(3):87-91.
50. Agematsu H, Watanabe H, Yamamoto H, Fukayama M, Kanazawa T, Miake K. Scanning electron microscopic observations of microcanals and continuous zones of interglobular dentin in human deciduous incisal dentin. *Bull Tokyo Dent Coll* 1990;31(2):163-73.
51. Sumikawa DA, Marshall GW, Gee L, Marshall SJ. Microstructure of primary tooth dentin. *Pediatr Dent* 1999;21(7):439-44.
52. Agematsu H, Sawada T, Watanabe H, Yanagisawa T, Ide Y. Immuno-scanning electron microscope characterization of large tubules in human deciduous dentin. *Anat Rec* 1997;248(7):339-45.
53. Dyngeland T, Fosse G. Scanning electron microscopic, light microscopic and micro-

- radiographic study of giant tubules in bovine dentin. *Scand J Dent Res* 1986;94(4):285-98.
54. Vujosevic L, Obradovic-Duricic K. Porosity of hard dental tissues. *Stomatol Glas Srb* 1989;36(2):95-100.
55. Buurma B, Gu K, Rutherford RB. Transplantation of human pulpal and gingival fibroblasts attached to synthetic scaffolds. *Eur J Oral Sci* 1999;107(4):282-9.
56. Luder HU, Zappa U. Nature and attachment of cementum formed under guided conditions in human teeth: an electron microscopic study. *J Periodontol* 1998;69(8):889-98.
57. Adriaens PA, Edwards CA, De Boever JA, Loesche WJ. Ultrastructural observations on bacterial invasion in cementum and radicular dentin of periodontally diseased human teeth. *J Periodontol* 1988;59(8):493-503.
58. Frank RM. Ultrastructure of human dentine 40 years ago: progress and perspectives. *Arch Oral Biol* 1999;44(12):979-84.
59. Murray PE, About I, Lumley PJ, Franquin JC, Remusat M, Smith AJ. Human odontoblast cell numbers after dental injury. *J Dent* 2000;28(4):277-85.
60. Jontell M, Gunraj MN, Bergenholtz G. Immunocompetent cells in the normal dental pulp. *J Dent Res* 1987;66(6):1149-53.
61. Hamilton JG. Needle phobia: a neglected diagnosis. *J Fam Pract* 1995;41(2):169-75.
62. Bedrock RD, Skigen A, Dolwick MF. Retrieval of a broken needle in the pterygo-mandibular space. *JADA* 1999;130(5):685-7.
63. Kuboki Y, Sasaki M, Saito A, Takita H, Kato H. Regeneration of periodontal ligament and cementum by BMP-applied tissue engineering. *Eur J Oral Sci* 1998;106(suppl 1):197-203.
64. Ripamonti U, Reddi AH. Tissue engineering, morphogenesis, and regeneration of the periodontal tissues by bone morphogenetic proteins. *Crit Rev Oral Biol Med* 1997;8(2):154-63.
65. Ripamonti U, Heliotis M, Rueger DC, Sampath TK. Induction of cementogenesis by recombinant human osteogenic protein-1 (hop-1/bmp-7) in the baboon (*Papio ursinus*). *Arch Oral Biol* 1996;41(1):121-6.
66. Ziccardi VB, Buchbinder D. Guided tissue regeneration in dentistry. *N Y State Dent J* 1996;62(10):48-51.
67. Cochran DL, Wozney JM. Biological mediators for periodontal regeneration. *Periodontol* 2000 1999;19(February):40-58.
68. Charette MF, Rutherford RB. Regeneration of dentin. In: Lanza RP, Langer RS, Chick WL, eds. Principles of tissue engineering. Austin, Texas: R.G. Landes; 1997:727-34.
69. Bohl KS, Shon J, Rutherford B, Mooney DJ. Role of synthetic extracellular matrix in development of engineered dental pulp. *J Biomater Sci Polym Ed* 1998;9(7):749-64.
70. Somerman MJ, Ouyang HJ, Berry JE, et al. Evolution of periodontal regeneration: from the roots' point of view. *J Periodontol Res* 1999;34(7):420-4.
71. Jeon KW, Lorch IJ, Danielli JF. Reassembly of living cells from dissociated components. *Science* 1970;167:1626-7.
72. Pressman EK, Levin IM, Sandakhchiev LS. Reassembly of an *Acetabularia mediterranea* cell from the nucleus, cytoplasm, and cell wall. *Protoplasma* 1973;76(3):327-32.
73. Morowitz HJ. Manufacturing a living

- organism. *Hosp Pract* 1974;9(11):210-5.
74. ADA Council on Scientific Affairs. Dental amalgam: update on safety concerns *JADA* 1998;129:494-503.
75. Absi EG, Addy M, Adams D. Dental hypersensitivity: a study of the patency of dentinal tubules in sensitive and non-sensitive cervical dentine. *J Clin Periodontol* 1987;14(5):280-4.
76. Addy M, West N. Etiology, mechanisms, and management of dentine hypersensitivity. *Curr Opin Periodontol* 1994;2:71-7.
77. Shellhart WC, Oesterle LJ. Uprighting molars without extrusion. *JADA* 1999;130:381-5.
78. Fartash B, Tangerud T, Silness J, Arvidson K. Rehabilitation of mandibular edentulism by single crystal sapphire implants and overdentures: 3-12 year results in 86 patients—a dual center international study. *Clin Oral Implants Res* 1996;7(3):220-9.
79. Castelnovo J, Tjan AH, Phillips K, Nicholls JI, Koic JC. Fracture load and mode of failure of ceramic veneers with different preparations. *J Prosthet Dent* 2000;83(2):171-80.
80. Fartash B, Hultin M, Gustafsson A, Asman B, Arvidson K. Markers of inflammation in crevicular fluid from peri-implant mucosa surrounding single crystal sapphire implants. *Clin. Oral Implants Res* 1997;8(1):32-8.
81. Wang ZX, Chai BF, Ye YQ, Fang QY. Local changes in aluminium, calcium and phosphorus content of bone caused by aluminium implant. *Chin Med J (Engl)* 1992;105(9):749-52.
82. Blankenau R, Goldstein RE, Haywood VB. The current status of vital tooth whitening techniques. *Compend Contin Educ Dent* 1999;20(8):781-9.
83. Kumar JV, Siegal MD. A contemporary perspective on dental sealants. *J Calif Dent Assoc* 1998;26(5):378-85.
84. Piattelli A, Podda G, Scarano A. Histological evaluation of bone reactions to aluminum oxide dental implants in man: a case report. *Biomaterials* 1996;17(7):711-4.
85. Kon M, Kuwayama N. Effect of adding diamond particles on the fracture toughness of apatite ceramics. *Dent Mater J* 1993;12(1):12-7.
86. Iijima S, Brabec C, Maiti A, Bernholc J. Structural flexibility of carbon nanotubes. *J Chem Phys* 1996;104:2089-92.
87. Brown LJ, Wall TP, Lazar V. Trends in total caries experience: permanent and primary teeth. *JADA* 2000;131:223-31.
88. Hajishengallis G, Michalek SM. Current status of a mucosal vaccine against dental caries. *Oral Microbiol Immunol* 1999;14(1):1-20.
89. Keller O, Orland FJ, Baird G. Ultrastructure of entamoeba gingivalis. *J Dent Res* 1967;46(5):1010-8.
90. Kleinberg I, Codipilly M. Modeling of the oral malodor system and methods of analysis. *Quintessence Int* 1999;30(5):357-69.
91. Roco MC. National nanotechnology initiative. Available at: "www.nano.gov". Accessed Sept. 26, 2000.
92. Zyvex Corp. Web site. Available at: "www.zyvex.com". Accessed Sept. 26, 2000.