



Supplemental material is available online.

Benefits and harms associated with analgesic medications used in the management of acute dental pain

An overview of systematic reviews

Paul A. Moore, DMD, PhD, MPH; Kathleen M. Ziegler, PharmD; Ruth D. Lipman, PhD; Anita Aminoshariae, DDS, MS; Alonso Carrasco-Labra, DDS, MSc; Angelo Mariotti, DDS, PhD

ABSTRACT

Background. Effective pain management is a priority in dental practice. Government and private agencies highlight the need to provide optimal pain relief, balancing potential benefits and harms of both opioid and nonopioid analgesic agents. The purpose of this study is to summarize the available evidence on the benefits and harms of analgesic agents, focusing on preexisting systematic reviews.

Types of Studies Reviewed. An overview of systematic reviews was conducted to evaluate the efficacy or reported adverse events associated with orally administered medication or medication combinations for relief of acute pain. Reviews were inclusive of all age populations but were limited to those that evaluated medication and medication combinations marketed in the United States and had moderate or high methodological quality according to the A MeaSurement Tool to Assess systematic Reviews (AMSTAR) 2 tool.

Results. Five reviews were found eligible for inclusion. The data identified combinations of ibuprofen and acetaminophen as having the highest association with treatment benefit in adult patients and the highest proportion of adult patients who experienced maximum pain relief. Diflunisal, acetaminophen, and oxycodone were found to have the longest duration of action in adult patients. Medication and medication combinations that included opioids were among those associated most frequently with acute adverse events in both child and adult-aged patient populations.

Practical Implications. The best available data suggested that the use of nonsteroidal medications, with or without acetaminophen, offered the most favorable balance between benefits and harms, optimizing efficacy while minimizing acute adverse events.

Key Words. Analgesia; pain relief; adverse events; systematic review; decision-making opioids; nonsteroidal anti-inflammatory drugs; opioids; acetaminophen.

JADA 2018;149(4):256-268

<https://doi.org/10.1016/j.adaj.2018.02.012>



Safe and effective pain management is an essential goal for all dental practitioners. Oral formulations, including both opioid and nonopioid analgesic agents, are among the medications commonly provided to manage pain for dental patients. Although the 2016 recommendations from the Centers for Disease Control and Prevention (CDC) about management of long-term pain were less well codified with respect to analgesic use for acute pain, it did include recommendations about limiting dose and duration of opioid-containing medications.¹ This likely reflects growing concern about the increasing occurrence of opioid misuse leading to deaths from both prescription and illegal opioids.² Although effective redress of this problem will be multifaceted, it will likely result in increased scrutiny about the choice of medications to be used when managing acute pain. The dual goals for pain management are safety and efficacy.

A variety of oral formulations of prescription and over-the-counter analgesic agents are often included alone or in combination as a component in strategies to manage acute dental pain.³

Copyright © 2018 American Dental Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Clinical considerations when determining the analgesic agent to be used include, but are not limited to, severity of the pain, patient pain sensitivity, medical history, specific dental pathologic process, and, in postoperative situations, the degree of surgical trauma.

The objective of our study is to summarize the data on oral analgesic medications with the aim of creating a compendium that details both the benefits and harms of these medications as a resource for dentists to use in their clinical decision making. This work was conducted in response to a request from the American Dental Association (ADA) Council on Dental Practice, using a protocol established a priori (available from the authors) and registered with the International Prospective Register of Systematic Reviews (PROSPERO) database (no. CRD42017080270) to summarize the best available evidence with respect to questions of safety and efficacy for relief of acute pain relevant to dental practice in the United States.

METHODS

This overview of reviews used the rapid review methodology⁴ to identify and summarize the available evidence from existing systematic reviews that examined the relative safety and efficacy of oral opioid and nonopioid analgesic agents available for use in the United States for the management of acute postoperative dental pain.

Selection criteria of included reviews

Type of Studies

We included systematic reviews and overviews of reviews with or without meta-analysis. We considered a report to be a systematic review by using a combination of selection criteria

- identified by the authors as a systematic review;
- included an explicit description of the search strategy;
- conducted the search in at least 2 electronic databases.

In addition, we selected reviews that ranked as moderate to high methodological quality according to the A MeaSurement Tool to Assess systematic Reviews (AMSTAR) 2 tool.⁵ Narrative reviews, editorials, and letters to the editor were excluded.

Type of Participants and Interventions

Systematic reviews or overviews of reviews that summarized data on the use of orally administered medications for the management of acute pain from studies that involved either adults or children were eligible for inclusion. The source of pain was mostly acute postoperative dental pain (for example, following third-molar extraction).

Type of Outcome Measures

Systematic reviews or overviews of reviews with data on the pharmacologic management of acute pain that reported on efficacy of pain relief (defined as at least 50% relief from maximum pain that lasted 46 hours), duration of pain relief (time before rescue remediation was requested), or any acute adverse events were included in this review.

Search methods for systematic review retrieval

The literature search strategy used the key words “(acute pain) AND (dental OR dentist* OR postop* OR postsurg*)” and was performed with the PubMed Clinical Queries for Systematic Reviews tool on April 13, 2017. In addition, manual searches of the reference lists of key articles were conducted to complement the electronic search. We also searched PROSPERO to identify systematic reviews under development that may have been relevant for our study.

Study selection, data collection, and analysis

The preliminary screening of titles and abstracts for all potentially eligible citations identified in the literature search was conducted in duplicate with the use of EndNote (Clarivate Analytics). In a second stage, the full text of any citation considered as potentially eligible was retrieved, and the eligibility was assessed. In case of disagreements among screeners, a third researcher acted as arbiter.

Assessment of the Methodological Quality of Reviews

We used the AMSTAR 2 tool to evaluate the methodological quality of the potentially eligible systematic reviews,⁵ and we used the AMSTAR rating of the individual reviews included in the

ABBREVIATION KEY

ADA:	American Dental Association.
AMSTAR:	A MeaSurement Tool to Assess systematic Reviews.
CDC:	Centers for Disease Control and Prevention.
NNTB:	Number needed to treat for benefit.
NNTH:	Number needed to treat to harm.
PROSPERO:	International Prospective Register of Systematic Reviews.




Figure. Scheme of citations identified in the PubMed Clinical Queries search.

print & web 4C/FPO

overviews of reviews provided by the overview authors. The AMSTAR 2 tool includes 16 items; only reviews that did not contain critical flaws were considered to provide a sufficiently accurate summary of the results from available studies to be included in our overview.

Data Extraction

After identification of eligible reviews, we collected numerical and narrative data reporting on desirable and undesirable outcomes when using any type of analgesic medication. The data were extracted by 1 researcher and entered directly into tables that were checked for accuracy by a second researcher in an independent fashion. We used the DynaMed Plus database (EBSCO Information Services) to determine which medications or medication combinations were marketed in the United States so as to only report data on medications or medication combinations marketed as of May 2017 in the United States.

Absolute Estimates to Summarize Study Results

We present analgesic efficacy data as the number of patients needed to treat for benefit (NNTB) along with the 95% confidence interval (CI) and the proportion of people attaining at least 50% maximum pain relief over 4 to 6 hours. In terms of pain medications, the NNTB is the number of people who must be treated by a specific dose of pain medication to receive 50% pain relief, which is considered clinically meaningful pain relief.⁶ The lower the NNTB, the more effective the analgesic agent. For example, an analgesic agent with an NNTB of 1 means that the medicine is 100% effective at reducing pain by 50%; that is, everyone who takes the medicine has effective pain relief. A drug with an NNTB of 2 means that 2 people must be treated for 1 to receive effective relief. According to a National Safety Council report,⁷ for oral pain medications, an NNTB of 1.5 would be considered very good and an NNTB of 2.5 would be considered good, whereas a drug with an NNTB of 10 would not be considered an effective analgesic (that is, 10 patients would have to be treated for 1 patient to experience pain relief). Analogous to NNTB, to compare risk of harm, the number needed to treat to harm (NNTH) can be calculated. NNTH is the number of patients needed to be treated for an additional adverse event to occur compared with the placebo group.

RESULTS

No ongoing or completed systematic reviews were identified through the search of the PROSPERO database that were relevant to this work. The PubMed Clinical Queries search identified 556 citations (Figure). Title and abstract screening of the retrieved citations, as well as any identified in manual searches of the reference lists of key articles, narrowed the results set of potential citations to 105.

Of these, a total of 74 (Appendix 1, available online at the end of this article) were excluded from the evidence review for having been superseded by subsequent systematic reviews or meta-analysis; 1 was excluded for not answering a relevant question; 4 were excluded for reviewing oral agents; and 6 were excluded for not being systematic reviews. After a review of the remaining 20 citations at the full-text level,⁸⁻²⁷ 15 were excluded (Appendix 2, available online at the end of this article) because they were included in subsequent reviews, were not systematic reviews, included studies that involved preoperative medication, examined an intervention unlikely to have a pharmacologic basis for efficacy, or reviewed medications that were not marketed in the United States. The remaining 5 systematic reviews or overviews of reviews were included.^{17,18,21,22,24} Two of these specifically targeted pediatric populations; 1 compared as-required versus fixed-schedule dosing of analgesic agents for postoperative pain in children,¹⁷ and the other focused on the safety of common analgesic agents for acute pain (albeit, nonsurgical) in the pediatric population.¹⁸

Among the systematic reviews that met the inclusion criteria, the 3 that explored pain management questions in adult populations^{21,22,24} explicitly indicated that they used AMSTAR to assess methodological quality.²⁸ Of the 2 systematic reviews that explored pain management in pediatric populations, 1¹⁷ simply documented adherence to these same principles for assessing the methodological quality of the systematic reviews they included, and the other¹⁸ reported using the McMaster Quality Assessment Scale of Harms.²⁹ Along with appraising the literature on which they were based, 4 of the systematic reviews^{17,21,22,24} also satisfied the AMSTAR 2 criteria for high methodological quality.⁵ Appraisal of the work by Hartling and colleagues¹⁸ found it to be of moderate methodological quality using AMSTAR 2.⁵

Relief of postoperative pain with the use of pharmacologic agents in adults

The data from randomized controlled trials that studied single-dose oral analgesic agents in acute postoperative pain come almost exclusively from studies that involved people after extraction of third molars. It derives from an overview of reviews of 39 Cochrane reviews by Moore and colleagues,²² which included only randomized clinical trials. The results are from more than 58,000 adult participants (15 years or older) in approximately 460 individual studies.

Evidence was considered high quality if data were available from at least 2 studies, each of which included at least 200 participants, and the results were of low risk of publication bias.²³ High-quality evidence was available for 53 medication and medication combinations. Several studied more than 1 dose in “painful postsurgical conditions”; these included various fixed-dose combinations and fast-acting formulations of some analgesic agents. Table 1 provides data for drugs available in the United States with the use of NNTB in order from most to least effective. NNTB ranged from 1.5 to 12 for at least 50% maximum pain relief over 4 to 6 hours compared with placebo with the proportion of participants achieving this level of benefit ranging from approximately 77% to 26% compared with 40% to 0% for placebo.

Medication and medication combinations with the lowest NNTBs, meaning that patients were most likely to experience treatment benefit, were 400 milligrams of ibuprofen plus 1,000 mg of acetaminophen with an NNTB of 1.5 (95% CI, 1.4 to 1.7), 200 mg of ibuprofen plus 500 mg of acetaminophen with an NNTB of 1.6 (95% CI, 1.5 to 1.8), 1,000 mg of acetaminophen plus 10 mg of oxycodone with an NNTB of 1.8 (95% CI, 1.6 to 2.2), and 100 mg of diclofenac potassium with an NNTB of 1.9 (95% CI, 1.7 to 2.3).

Medication and medication combinations with the highest proportion of patients who experienced at least 50% maximum pain relief for 4 to 6 hours were 600 mg of ibuprofen (77%), 400 mg of ibuprofen plus 1,000 mg of acetaminophen (72%), 200 mg of ibuprofen plus 500 mg of acetaminophen (69%), and 50 mg of flurbiprofen (69%) (Table 1).

Duration of postoperative pain relief with the use of pharmacologic agents in adults

Duration of postoperative pain relief, assessed as the time to remedication ranged from 1.5 hours (placebo) to more than 20 hours (Table 1).²² Medication and medication combinations with the longest duration of action, meaning that they had the longest time before rescue remedication was requested, was 10.9 hours for 1,000 mg of diflunisal, 9.9 hours for 650 mg of acetaminophen and 10 mg of oxycodone, 9.8 hours for 500 mg of diflunisal, and 8.9 hours for 500 to 550 mg of naproxen. The medications and medication combinations that provided the shortest duration of pain

Table 1. Efficacy data from high-quality studies for analgesic agents available in the United States in order of effectiveness (most to least) according to NNTB*.^{22,24}

DRUG OR DRUG COMBINATION, DOSE	NNTB	95% CONFIDENCE INTERVAL	AT LEAST 50% MAXIMUM PAIN RELIEF OVER 4-6 HOURS, %		MEAN OR MEDIAN TIME TO REMEDIATION, HOURS	
			Active	Placebo	Active	Placebo
Ibuprofen Plus Acetaminophen, 400 Milligrams/1,000 mg	1.5	1.4 to 1.7	72	6	8.3	1.7
Ibuprofen Plus Acetaminophen, 200 mg/500 mg	1.6	1.5 to 1.8	69	6	7.6	1.7
Acetaminophen Plus Oxycodone, 1,000 mg/10 mg	1.8	1.6 to 2.2	68	13	9.8	1.5
Diclofenac (Potassium), 100 mg	1.9	1.7 to 2.3	65	13	6.3	2.0
Ketoprofen, 25 mg	2.0	1.8 to 2.3	62	12	46 [†]	79 [†]
Diclofenac (Potassium), 50 mg	2.1	1.9 to 2.5	64	17	4.5	1.7
Diflunisal, 1,000 mg	2.1	1.8 to 2.6	62	15	10.9	3.2
Ibuprofen (Fast-Acting), 200 mg	2.1	1.9 to 2.4	57	10	43 [†]	78 [†]
Ibuprofen (Fast-Acting), 400 mg	2.1	1.9 to 2.3	65	18	32 [†]	82 [†]
Ibuprofen Plus Caffeine, 100 mg/200 mg	2.1	1.9 to 3.1	59	10	26 [†]	60 [†]
Ketoprofen, 100 mg	2.1	1.7 to 2.6	66	18	43 [†]	85 [†]
Acetaminophen Plus Codeine, 800-1,000 mg/60 mg	2.2	1.8 to 2.9	53	7	5.0	2.3
Ibuprofen Plus Codeine, 400 mg/26-60 mg	2.2	1.8 to 2.6	64	18	NA [§]	
Fenoprofen, 200 mg	2.3	1.9 to 3.0	57	13	NA	
Ibuprofen Plus Oxycodone, 400 mg/10 mg	2.3	2.0 to 2.8	60	17	NA	
Aspirin, 1,200 mg	2.4	1.9 to 3.2	62	19	NA	
Diclofenac (Fast-Acting), 50 mg	2.4	2.0 to 3.0	61	20	7.6	3.8
Diclofenac (Potassium), 25 mg	2.4	2.0 to 2.9	56	15	3.1	1.2
Ibuprofen Plus Caffeine, 100 mg/100 mg	2.4	1.9 to 3.1	43	0	34 [†]	79 [†]
Ketoprofen, 12.5 mg	2.4	1.9 to 3.1	56	13	80 [†]	98 [†]
Flurbiprofen, 100 mg	2.5	2.0 to 3.1	65	24	16 [†]	68 [†]
Ibuprofen (Acid), 400 mg	2.5	2.4 to 2.6	52	12	5.6	1.9
Celecoxib, 400 mg	2.6	2.3 to 3.0	43	5	8.4	1.6
Diflunisal, 500 mg	2.6	2.1 to 3.3	53	14	9.8	3.2
Acetaminophen Plus Oxycodone, 650 mg/10 mg	2.7	2.4 to 3.1	51	14	9.8	1.5
Flurbiprofen, 50 mg	2.7	2.3 to 3.3	69	32	25 [†]	66 [†]
Ibuprofen (Acid), 600 mg	2.7	2.0 to 4.2	77	40	NA	
Naproxen, 400-440 mg	2.7	2.2 to 3.5	49	11	NA	
Naproxen, 500-550 mg	2.7	2.3 to 3.3	52	15	8.9	2.0
Piroxicam, 20 mg	2.7	2.1 to 3.8	63	26	NA	
Etodolac, 400 mg	2.9	2.3 to 4.0	39	5	NA	
Ibuprofen (Acid), 200 mg	2.9	2.7 to 3.2	41	7	4.7	2.1
Etodolac, 200 mg	3.3	2.7 to 4.2	44	13	NA	
Flurbiprofen, 25 mg	3.3	2.5 to 4.9	35	5	35 [†]	70 [†]
Ketoprofen, 50 mg	3.3	2.7 to 4.3	48	18	48 [†]	81 [†]
Acetaminophen, 500 mg	3.5	2.7 to 4.8	61	32	35 [†]	63 [†]
Acetaminophen, 975-1,000 mg	3.6	3.2 to 4.1	46	18	3.9	2.7
Acetaminophen Plus Codeine, 600-650 mg/60 mg	3.9	3.3 to 4.7	43	17	4.1	2.4
Aspirin, 600-650 mg	4.2	3.8 to 4.6	39	15	55 [†]	75 [†]
Aspirin, 1,000 mg	4.2	3.8 to 4.6	41	14	67 [†]	83 [†]

* NNTB: Number needed to treat for benefit. † Percentage remediating within 6 hours. ‡ Percentage remediating within 8 hours. § NA: Not available.

Table 1. Continued

DRUG OR DRUG COMBINATION, DOSE	NNTB	95% CONFIDENCE INTERVAL	AT LEAST 50% MAXIMUM PAIN RELIEF OVER 4-6 HOURS, %		MEAN OR MEDIAN TIME TO REMEDIATION, HOURS	
			Active	Placebo	Active	Placebo
Celecoxib, 200 mg	4.2	3.4 to 5.6	35	11	6.6	2.6
Ibuprofen (Acid), 100 mg	4.3	3.2 to 6.4	31	8	NA	
Acetaminophen, 600-650 mg	4.6	3.9 to 5.5	38	16	3.5	2.4
Etodolac, 100 mg	4.8	3.5 to 7.8	41	20	NA	
Gabapentin, 250 Milliliters	11.0	6.4 to 35	15	5	2.4	2.1
Codeine, 60 mg	12.0	8.4 to 18	26	17	2.7	2.0

Table 2. Acute adverse events for medication or medication combination with statistically significant difference from placebo controls.²¹

DRUG OR DRUG COMBINATION, DOSE	NUMBER NEEDED TO TREAT TO HARM	95% CONFIDENCE INTERVAL	REPORTED ACUTE ADVERSE EVENT, %	
			Active	Placebo
Ibuprofen Plus Caffeine, 200 Milligrams/100 mg	2.2	1.0 to 4.9	11	6
Acetaminophen Plus Oxycodone, 650 mg/10 mg	1.8	1.4 to 2.3	58	28
Diflunisal, 1,000 mg	1.8	1.2 to 2.6	29	16
Acetaminophen Plus Codeine, 600-650 mg/60 mg	1.6	1.3 to 1.9	34	17
Acetaminophen Plus Oxycodone, 1,000 mg/10 mg	1.6	1.3 to 2.0	68	43
Aspirin, 1,000 mg	1.6	1.1 to 2.3	26	12
Acetaminophen Plus Codeine, 800-1,000 mg/60 mg	1.4	1.2 to 1.6	31	19
Ibuprofen Plus Acetaminophen, 200 mg/500 mg*	0.7	0.6 to 0.9	30	48
Ibuprofen Plus Acetaminophen, 400 mg/1,000 mg*	0.6	0.5 to 0.8	29	48

* Statistically fewer adverse events in medication combination than in placebo controls.

relief were 3.5 hours for 600 to 650 mg of acetaminophen, 3.1 hours for 25 mg of diclofenac potassium, 2.7 hours for 60 mg of codeine, and 2.4 hours for 250 mg of gabapentin.

Adverse events associated with provision of pharmacologic agents for relief of postoperative pain

Studies of medications for relief of acute pain are designed and powered for primary outcomes that have to do with analgesia rather than adverse events. Although reporting of adverse events in clinical trials is generally poor and variable depending on methodology, Moore and colleagues²¹ have summarized the data on adverse events available from studies of sufficient quality to have been included in Cochrane reviews of pharmacologic agents used for single-dose oral analgesic agents for acute postoperative pain.^{22,24} Although noting that accurate estimation of frequency or severity of adverse events is difficult at best, they calculated that serious, acute adverse events are rare and estimated that they occur in approximately 1 in 3,200 people.²¹

Methodologically, information about acute adverse events was most often gathered through the use of patient diaries. Adverse events included drowsiness, respiratory depression, nausea, vomiting, and constipation for the opioid medications and drowsiness, dizziness, nausea, and headache for the nonsteroidal anti-inflammatory drugs, although there was no effort made to parse the frequency of specific adverse events.²¹ Table 2 presents the relatively short list of medication and medication combinations with a statistically significant difference in the proportion of patients who reported an adverse event compared with patients in the placebo group. Although the largest NNTH reported was for 200 mg of ibuprofen and 100 mg of caffeine (2.2; 95% CI, 1.0 to 4.9), 4 of the 6 other

Table 3. Alphabetical listing of medication and medication combination with no statistically significant difference in acute adverse events compared with placebo controls.²¹

DRUG OR DRUG COMBINATION, DOSE	NNT ^H	95% CONFIDENCE INTERVAL	REPORTED ACUTE ADVERSE EVENT, %	
			Active	Placebo
Acetaminophen, 500 milligrams	0.9	0.4 to 1.9	7	6
Acetaminophen, 600-650 mg	1.2	0.9 to 1.5	16	14
Acetaminophen, 975-1,000 mg	1.1	0.9 to 1.3	18	16
Aspirin, 600-650 mg	1.2	1.0 to 1.4	11.0	9.5
Codeine, 60 mg	1.3	0.9 to 1.7	20	16
Diclofenac (Fast-Acting), All Doses	1.0	0.6 to 1.8	8	46
Diclofenac (Potassium), All Doses	1.0	0.7 to 1.6	8	46
Diflunisal, 500 mg	1.3	0.8 to 1.9	18	15
Etodolac, 100 mg	1.6	0.9 to 2.8	11	7
Etodolac, 200 mg	1.2	0.9 to 1.7	22	17
Etodolac, 400 mg	0.8	0.5 to 1.2	28	34
Fenoprofen, 200 mg	0.9	0.4 to 2.1	6	6
Flurbiprofen, 100 mg	1.0	0.6 to 1.8	12	12
Flurbiprofen, 25 mg	0.9	0.5 to 1.7	14	16
Flurbiprofen, 50 mg	0.8	0.5 to 1.1	13	17
Gabapentin, 250 mg	0.9	0.7 to 1.3	28	32
Ibuprofen, 50 mg	1.3	0.6 to 3.0	10	7
Ibuprofen, 100 mg	1.2	0.7 to 2.1	14	13
Ibuprofen, 200 mg	0.9	0.7 to 1.02	19	19
Ibuprofen, 400 mg	0.9	0.8 to 1.04	17	16
Ibuprofen Plus Caffeine, 100 mg/100 mg	1.9	0.8 to 4.1	14	8
Ibuprofen Plus Codeine, 400 mg/26-60 mg	1.2	0.8 to 1.7	28	19
Ibuprofen Plus Oxycodone, 400 mg/10 mg	1.2	0.8 to 1.7	28	19
Ketoprofen, 100 mg	1.2	0.7 to 2.2	22	18
Ketoprofen, 12.5 mg	1.3	0.5 to 3.6	6	4
Ketoprofen, 25 mg	1.2	0.7 to 2.0	10	10
Ketoprofen, 50 mg	1.6	0.9 to 2.6	21	14
Naproxen, 400-440 mg	1.3	0.8 to 2.2	22	17
Naproxen, 500-550 mg	1.0	0.7 to 1.2	27	29
Oxycodone, 5 mg	1.1	0.8 to 1.6	31	29

medication and medication combinations in which the NNT^H was statistically greater than placebo were for opioid-containing combinations (Tables 2 and 3). That the 2 combinations of ibuprofen and acetaminophen, at dosages of 200 mg/500 mg and 400 mg/1,000 mg, were observed to have statistically fewer adverse events in the active treatment group than in the placebo control group (Table 2) may reflect a sample size issue. The data on adverse events for the medication and medication combinations for which there was no statistically significant difference compared with the placebo control are presented in Table 3 in alphabetical order.

Relief of postoperative pain with the use of pharmacologic agents in children

A 2015 Cochrane review attempted to evaluate the management of postoperative pain in children younger than 16 years.¹⁷ Three randomized clinical trials that met their inclusion criteria were found, enrolling a total of 246 children. (The only medication for which the researchers found data

Table 4. Acute adverse events observed in children for medications or medication combinations.¹⁸

MEDICATION OR MEDICATION COMBINATION, DOSE	STUDIES, NO.	ACUTE ADVERSE EVENTS REPORTED, NO.	STUDY PARTICIPANTS, NO.	ACUTE ADVERSE EVENTS, %
Codeine, 2 milligrams/kilograms	1	74	56	132
Oxycodone, 0.2 mg/kg	2	69	73	95
Morphine, 0.5 mg/kg	2	84	140	60
Ibuprofen, 10 mg/kg, and Oxycodone, 0.1 mg/kg	1	8	22	36
Acetaminophen Plus Codeine, 1 mg/kg	3	94	368	26
Ibuprofen, 10 mg/kg, and Codeine, 1 mg/kg	2	52	209	25
Ibuprofen, 10 mg/kg	9	74	510	15
Naproxen, 20 mg/kg	1	4	41	10
Ketoprofen, 40 mg	1	2	33	6
Tramadol, 2 mg/kg	1	3	67	4
Acetaminophen, 15 mg/kg	6	10	260	4

sufficient for calculating the risk ratio with 95% CI was dipyrrone, which is only available for human use in Europe and Latin America. It is included here for completeness in reporting the results from this Cochrane review.) The researchers found moderate quality data to indicate that 500 mg of oral dipyrrone provided 70% of patients with at least 50% relief of pain at 4 to 6 hours (95% CI, 1.8 to 3.1). They reported being unable to conduct analysis of adverse events because data were inconsistently reported.

Adverse events associated with provision of pharmacologic agents for management of acute, nonsurgical pain in children

A systematic review by Hartling and colleagues¹⁸ on adverse events on children (defined as younger than 18 years) who received pharmacologic agents for managing acute pain included nausea; vomiting; headache; gastrointestinal symptoms other than nausea and vomiting; drowsiness, sleepiness, tiredness; dizziness; itchiness, rash, pruritus; central nervous system symptoms; and pulmonary symptoms. Although original data were insufficient to calculate NNTH, we qualitatively summarized the results in Table 4 which show that the medication and medication combinations that include opioids are associated with the largest proportion of acute adverse events, ranging from 132% (that is, more than 1 type of acute adverse event per patient) for 2 mg/kg of codeine, 95% for 0.2 mg/kg of oxycodone, and 60% for 0.5 mg/kg of morphine and, at the low end of the spectrum, 15% or fewer patients who reported an acute adverse event for 10 mg/kg of ibuprofen, 20 mg/kg of naproxen, 40 mg/kg of ketoprofen, 2 mg/kg of tramadol, or 15 mg/kg of acetaminophen. (The list of the medication and medication combinations are presented in descending order of study participants affected by adverse events in Table 4.)

DISCUSSION

To determine which oral analgesic medications to use for relief of acute dental pain appropriate for the patient, health care professionals should consider both the medication's potential to provide pain relief and its potential to cause harm. A variety of medication and medication combinations, including formulations that contain opioids, may be considered for the management of acute dental pain, and it is important to be cognizant that no medication or medication combination produces high levels of pain relief in all patients and that the analgesic agents prescribed are not intended to eliminate all pain that may present. When prescribing analgesic agents, practitioners should appreciate and counsel patients that the goal is for the patient to be as comfortable as possible, although patients should be aware that some discomfort is normal and may still occur. The range of results with single-dose analgesic agents in participants with moderate or severe acute pain was from 7 of 10 (70%) achieving good pain relief with the most effective medicine to approximately 3 of 10

(30%) with the least effective medicine. In terms of the decision-making process about what medication or medication combination to prescribe, the Joint Commission's statement on pain management indicates that pain management strategies should reflect a patient-centered approach and consider the patient's current presentation, the health care providers' clinical judgment, and the risks and benefits associated with the strategies, including potential risk of dependency, addiction, and abuse.³⁰ Although most data in adults presented here derive from the study of third-molar extraction, the results are more broadly applicable, because comparable findings, for example, have been reported for relief of pain of endodontic origin.³¹

When comparing the efficacy of nonsteroidal anti-inflammatory medications with opioids in relation to the magnitude of pain relief, the combination of 400 mg of ibuprofen plus 1,000 mg of acetaminophen was found to be superior to any opioid-containing medication or medication combination studied. In addition, the opioid-containing medications or medication combinations studied were all found to have higher risk of inducing acute adverse events than 400 mg of ibuprofen plus 1,000 mg of acetaminophen. Thus, in general, when considering either benefits or harms, management of acute pain with nonsteroidal medications, with or without acetaminophen, appears to have a therapeutic advantage to opioid-containing medications. Although there are situations in which clinical judgment indicates an opioid-containing medication may be warranted, the data make a compelling case favoring use of nonsteroidal medications, with or without acetaminophen.

Many factors contribute to prescribing decisions made by dentists, including education, training, and local legislation. There is reported geographic variation in opioid-prescribing patterns.³² No single common course curriculum, objectives, or assessments are used by all dental schools.³³ Another approach may be dental school and continuing education programs about the CDC guidelines for prescribing opioids for long-term pain,¹ which have been effective in changing opioid-prescribing patterns for clinicians in the fields of surgery³⁴ and emergency medicine.³⁵ Although data presented in this study do not cover the breadth of the CDC recommendations, they are relevant to the concept of optimizing nonopioid therapy before moving to a trial of opioids. This is consistent with the recent ADA Statement on the Use of Opioids in the Treatment of Dental Pain revised in October 2016, which indicates that "Dentists should consider nonsteroidal anti-inflammatory analgesics as the first-line therapy for acute pain management."³⁶

Strengths and limitations of current review

The AMSTAR 2 evaluations found that the information gleaned from the available systematic reviews on efficacy and acute adverse events is of moderate-high methodological quality. This should allow readers to have confidence that the use of appropriately selected nonsteroidal anti-inflammatory analgesic medication can deliver the desired pain management. Although studies are not designed with adverse events as a primary end point, with report of serious adverse events being minimal, clinicians can use this to extrapolate about the overall safety of these medications. In terms of limitations, the literature search strategy used was systematic, but broadening it to capture primary studies and eliminating the English-only language exclusion criterion may have yielded additional information. The data evaluated in randomized clinical trials to assess analgesic efficacy used a single-dose model of treatment and are unable to quantitatively evaluate adverse events that may require longer follow-up to be detected—that is, dependence or medication diversion that is increasingly seen to be problematic.

CONCLUSIONS

Opioid medication and medication combinations are not among the most effective or long lasting of the options available for relief of acute dental pain. In addition, opioid medication and medication combinations are associated with higher rates of acute adverse events. From the perspective of risk-benefit analysis, justifying general use of opioid medications as first-line therapy for management of acute pain remains unclear. The large set of published research reports summarized here suggests that relief of postoperative pain in dental practice with the use of nonsteroidal anti-inflammatory drugs, with or without acetaminophen, is equal or superior to that provided by opioid-containing medications. ■

SUPPLEMENTAL DATA

Supplemental data related to this article can be found at <https://doi.org/10.1016/j.adaj.2018.02.012>.

Dr. Moore is a professor in the Departments of Dental Anesthesiology and Dental Public Health, School of Dental Medicine, University of Pittsburgh, 386 Salk Hall, Pittsburgh, PA 15261, e-mail pam7@pitt.edu. Address correspondence to Dr. Moore.

Dr. Ziegler is a manager in Scientific Information, Science Institute, American Dental Association, Chicago, IL.

Dr. Lipman is the director of Scientific Information, Science Institute, American Dental Association, Chicago, IL.

Dr. Aminoshariae is an associate professor, Department of Endodontics, School of Dental Medicine, Case Western Reserve University, Cleveland, OH.

Dr. Carrasco-Labra is the director, Center for Evidence Based Dentistry, Science Institute, American Dental Association, Chicago, IL.

Dr. Mariotti is the chair, Division of Periodontology, College of Dentistry, The Ohio State University, Columbus, OH.

Disclosure. None of the authors reported any disclosures.

This work derives in part from a report requested by the American Dental Association (ADA) Council for Dental Practice as a resource for content development in relation to the ADA Statement on the Use of Opioids in the Treatment of Dental Pain adopted in October 2016 by the ADA House of Delegates.³⁵ The authors wish to acknowledge the ongoing efforts of the Council on Dental Practice and the Council on Government Affairs to raise member awareness about the complex and multifaceted issues having to do with medications prescribed or recommended for pain relief. The authors also thank Ms. Anita Mark, Senior Scientific Content Specialist, ADA Science Institute, for her valuable editorial contribution.

1. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain: United States, 2016. *MMWR Recomm Rep*. 2016;65(11):295.
2. Soelberg CD, Brown RE Jr, Du Vivier D, Meyer JE, Ramachandran BK. The US opioid crisis: current federal and state legal issues. *Anesth Analg*. 2017;125(5):1675-1681.
3. Ganzberg S, Friction J. Analgesics: opioids and nonopioids. In: Ciancio SG, ed. *ADA/PDR Guide to Dental Therapeutics*. 5th ed. Montvale, NJ: Physicians' Desk Reference; 2009:63-133.
4. Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J*. 2009;26(2):91-108.
5. Shea BJ, Reeves BC, Wells G, et al. AMSTAR 2: a critical appraisal tool for systematic reviews that include randomised or non-randomised studies of healthcare interventions, or both. *BMJ*. 2017;358:j4008.
6. Martin WJ, Ashton-James CE, Skorpil NE, Heymans MW, Forouzanfar T. What constitutes a clinically important pain reduction in patients after third molar surgery? *Pain Res Manag*. 2013;18(6):319-322.
7. Teater D. Evidence for the safety of pain medications. National Safety Council 2014. Available at: <http://www.nsc.org/RxDrugOverdoseDocuments/Evidence-Efficacy-Pain-Medications.pdf>. Accessed February 28, 2018.
8. American Academy of Pediatrics, Committee on Psychosocial Aspects of Child and Family Health; Task Force on Pain in Infants, Children, and Adolescents. The assessment and management of acute pain in infants, children, and adolescents. *Pediatrics*. 2001;108(3):793-797.
9. Alexander L, Hall E, Eriksson L, Rohlin M. The combination of non-selective NSAID 400 mg and paracetamol 1000 mg is more effective than each drug alone for treatment of acute pain: a systematic review. *Swed Dent J*. 2014;38(1):1-14.
10. Au AH, Choi SW, Cheung CW, Leung YY. The efficacy and clinical safety of various analgesic combinations for post-operative pain after third molar surgery: a systematic review and meta-analysis. *PLoS One*. 2015;10(6):e0127611.
11. Bu X, Yang L, Zuo Y. Efficacy and safety of perioperative parecoxib for acute postoperative pain treatment in children: a meta-analysis. *Front Med*. 2015;9(4):496-507.
12. Derry S, Cooper TE, Phillips T. Single fixed-dose oral dextropropofol plus tramadol for acute postoperative pain in adults. *Cochrane Database Syst Rev*. 2016;9:CD012232.
13. Derry S, Karlin SM, Moore RA. Single dose oral ibuprofen plus codeine for acute postoperative pain in adults. *Cochrane Database Syst Rev*. 2015;(2):CD010107.
14. Derry S, Wiffen PJ, Moore RA. Single dose oral ibuprofen plus caffeine for acute postoperative pain in adults. *Cochrane Database Syst Rev*. 2015;(7):CD011509.
15. Derry S, Wiffen PJ, Moore RA. Single dose oral diclofenac for acute postoperative pain in adults. *Cochrane Database Syst Rev*. 2015;(7):CD004768.
16. Eipe N, Penning J, Yazdi F, et al. Perioperative use of pregabalin for acute pain: a systematic review and meta-analysis. *Pain*. 2015;156(7):1284-1300.
17. Hobson A, Wiffen PJ, Conlon JA. As required versus fixed schedule analgesic administration for post-operative pain in children. *Cochrane Database Syst Rev*. 2015;(2):CD011404.
18. Hartling L, Ali S, Dryden DM, et al. How safe are common analgesics for the treatment of acute pain for children? A systematic review. *Pain Res Manag*. 2016;2016:5346819.
19. Hearn L, Derry S, Moore RA. Single dose dipyrone (metamizole) for acute postoperative pain in adults. *Cochrane Database Syst Rev*. 2016;4:CD011421.
20. Mishriky BM, Waldron NH, Habib AS. Impact of pregabalin on acute and persistent postoperative pain: a systematic review and meta-analysis. *Br J Anaesth*. 2015;114(1):10-31.
21. Moore RA, Derry S, Aldington D, Wiffen PJ. Adverse events associated with single dose oral analgesics for acute postoperative pain in adults: an overview of Cochrane reviews. *Cochrane Database Syst Rev*. 2015;(10):CD011407.
22. Moore RA, Derry S, Aldington D, Wiffen PJ. Single dose oral analgesics for acute postoperative pain in adults: an overview of Cochrane reviews. *Cochrane Database Syst Rev*. 2015;(9):CD008659.
23. Moore RA, Derry S, Straube S, Ireson-Paine J, Wiffen PJ. Faster, higher, stronger? Evidence for formulation and efficacy for ibuprofen in acute pain. *Pain*. 2014;155(1):14-21.
24. Moore RA, Wiffen PJ, Derry S, Maguire T, Roy RM, Tyrrell L. Non-prescription (OTC) oral analgesics for acute pain: an overview of Cochrane reviews. *Cochrane Database Syst Rev*. 2015;(11):CD010794.
25. Moore PA, Hersh EV. Combining ibuprofen and acetaminophen for acute pain management after third-molar extractions: translating clinical research to dental practice. *JADA*. 2013;144(8):898-908.
26. Nir RR, Nahman-Averbuch H, Moont R, Sprecher E, Yarnitsky D. Preoperative preemptive drug administration for acute postoperative pain: a systematic review and meta-analysis. *Eur J Pain*. 2016;20(7):1025-1043.
27. Raak C, Bussing A, Gassmann G, Boehm K, Ostermann T. A systematic review and meta-analysis on the use of *Hypericum perforatum* (St. John's Wort) for pain conditions in dental practice. *Homeopathy*. 2012;101(4):204-210.
28. Shea BJ, Grimshaw JM, Wells GA, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. *BMC Med Res Methodol*. 2007;7:10.
29. *Methods Guide for Effectiveness and Comparative Effectiveness Reviews*. Rockville, MD: Agency for Healthcare Research and Quality; 2008.
30. Baker DW. Joint Commission Statement on Pain Management. 2016. Available at: https://www.jointcommission.org/joint_commission_statement_on_pain_management/. Accessed February 28, 2018.
31. Aminoshariae A, Kulild JC, Donaldson M, Hersh EV. Evidence-based recommendations for analgesic efficacy to treat pain of endodontic origin: a systematic review of randomized controlled trials. *JADA*. 2016;147(10):826-839.
32. Paulozzi LJ, Mack KA, Hockenberry JM. Division of Unintentional Injury Prevention, National Center for Injury Prevention and Control, CDC. Vital signs: variation among States in prescribing of opioid pain relievers and benzodiazepines—United States, 2012. *MMWR Morb Mortal Wkly Rep*. 2014;63(26):563-568.
33. Clark DJ, Schumacher MA. America's opioid epidemic: supply and demand considerations. *Anesth Analg*. 2017;125(5):1667-1674.
34. Hill MV, Stucke RS, McMahon ML, Beeman JL, Barth RJ Jr. An educational intervention decreases opioid prescribing after general surgical operations. *Ann Surg*. 2018;267(3):468-472.
35. Osborn SR, Yu J, Williams B, Vasilyadis M, Blackmore CC. Changes in provider prescribing patterns after implementation of an emergency department prescription opioid policy. *J Emerg Med*. 2017;52(4):538-546.
36. American Dental Association Statement on the Use of Opioids in the Treatment of Dental Pain. 2016. Available at: <http://www.ada.org/en/about-the-ada/ada-positions-policies-and-statements/statement-on-opioids-dental-pain>. Accessed February 28, 2018.

APPENDIX 1

References excluded after title and abstract review

REASON: SUPERSEDED

- Barden J, Derry S, McQuay HJ, Moore RA. Single dose oral ketoprofen and dexketoprofen for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD007355.
- Barden J, Edwards J, Moore A, McQuay H. Single dose oral paracetamol (acetaminophen) for postoperative pain. *Cochrane Database Syst Rev.* 2004;(1):CD004602.
- Barden J, Edwards J, Moore RA, McQuay HJ. Single dose oral diclofenac for postoperative pain. *Cochrane Database Syst Rev.* 2004;(2):CD004768.
- Barden J, Edwards J, Moore RA, McQuay HJ. Single dose oral rofecoxib for postoperative pain. *Cochrane Database Syst Rev.* 2005;(1):CD004604.
- Barden J, Edwards J, Moore RA, McQuay HJ. Single dose oral rofecoxib for postoperative pain. *Cochrane Database Syst Rev.* 2004;(1):CD004604.
- Barden J, Edwards JE, McQuay HJ, Moore RA. Oral valdecoxib and injected parecoxib for acute postoperative pain: a quantitative systematic review. *BMC Anesthesiol.* 2003;3(1):1.
- Barden J, Edwards JE, McQuay HJ, Moore RA. Single dose oral celecoxib for postoperative pain. *Cochrane Database Syst Rev.* 2003;(2):CD004233.
- Barden J, Edwards JE, McQuay HJ, Moore RA. Single-dose rofecoxib for acute postoperative pain in adults: a quantitative systematic review. *BMC Anesthesiol.* 2002;2(1):4.
- Barden J, Edwards JE, McQuay HJ, Wiffen PJ, Moore RA. Relative efficacy of oral analgesics after third molar extraction. *Br Dent J.* 2004;197(7):407-411; discussion 397.
- Barreveld A, Witte J, Chahal H, Durieux ME, Strichartz G. Preventive analgesia by local anesthetics: the reduction of postoperative pain by peripheral nerve blocks and intravenous drugs. *Anesth Analg.* 2013;116(5):1141-1161.
- Bulley S, Derry S, Moore RA, McQuay HJ. Single dose oral rofecoxib for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD004604.
- Cicconetti A, Bartoli A, Ripari F, Ripari A. COX-2 selective inhibitors: a literature review of analgesic efficacy and safety in oral-maxillofacial surgery. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2004;97(2):139-146.
- Clarke R, Derry S, Moore RA, McQuay HJ. Single dose oral etoricoxib for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(2):CD004309.
- Clarke R, Derry S, Moore RA. Single dose oral etoricoxib for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2014;(5):CD004309.
- Clarke R, Derry S, Moore RA. Single dose oral etoricoxib for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2012;(4):CD004309.
- Derry C, Derry S, Moore RA, McQuay HJ. Single dose oral ibuprofen for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD001548.
- Derry C, Derry S, Moore RA, McQuay HJ. Single dose oral naproxen and naproxen sodium for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(1):CD004234.
- Derry CJ, Derry S, Moore RA. Single dose oral ibuprofen plus paracetamol (acetaminophen) for acute postoperative pain. *Cochrane Database Syst Rev.* 2013;(6):CD010210.
- Derry P, Derry S, Moore RA, McQuay HJ. Single dose oral diclofenac for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(2):CD004768.
- Derry S, Barden J, McQuay HJ, Moore RA. Single dose oral celecoxib for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2008;(4):CD004233.
- Derry S, Best J, Moore RA. Single dose oral dexibuprofen [S(+)-ibuprofen] for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2013;(10):CD007550.
- Derry S, Derry CJ, Moore RA. Single dose oral ibuprofen plus oxycodone for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2013;(6):CD010289.
- Derry S, Karlin SM, Moore RA. Single dose oral ibuprofen plus codeine for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2013;(3):CD010107.
- Derry S, Moore RA, McQuay HJ. Single dose oral codeine, as a single agent, for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2010;(4):CD008099.
- Derry S, Moore RA. Single dose oral aspirin for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2012;(4):CD002067.
- Derry S, Moore RA. Single dose oral celecoxib for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2013;(10):CD004233.
- Derry S, Moore RA. Single dose oral celecoxib for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2012;(3):CD004233.
- Dodson T. Paracetamol is an effective drug to use for pain following oral surgery. *Evid Based Dent.* 2007;8(3):79-80.
- Edwards J, Meseguer F, Faura C, et al. Single dose dipyron for acute postoperative pain. *Cochrane Database Syst Rev.* 2010;(9):CD003227.
- Elvir-Lazo OL, White PF. The role of multimodal analgesia in pain management after ambulatory surgery. *Curr Opin Anaesthesiol.* 2010;23(6):697-703.
- Francesca F, Bader P, Echte D, Giunta F, Williams J. EAU guidelines on pain management. *Eur Urol.* 2003;44(4):383-389.
- Gaskell H, Derry S, Moore RA, McQuay HJ. Single dose oral oxycodone and oxycodone plus paracetamol (acetaminophen) for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD002763.
- Hall PE, Derry S, Moore RA, McQuay HJ. Single dose oral lornoxicam for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD007441.
- Jordan RA, Hong HM, Lucaciu A, Zimmer S. Efficacy of straight versus angled interdental brushes on interproximal tooth cleaning: a randomized controlled trial. *Int J Dent Hyg.* 2014;12(2):152-157.
- Kakkar M, Derry S, Moore RA, McQuay HJ. Single dose oral nefopam for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007442.
- Mason L, Edwards J, Moore RA, McQuay HJ. Single dose oral indometacin for the treatment of acute postoperative pain. *Cochrane Database Syst Rev.* 2004;(4):CD004308.
- McQuay H, Edwards J. Meta-analysis of single dose oral tramadol plus acetaminophen in acute postoperative pain. *Eur J Anaesthesiol Suppl.* 2003;28:19-22.
- Moll R, Derry S, Moore RA, McQuay HJ. Single dose oral mefenamic acid for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2011;(3):CD007553.
- Moore OA, McIntyre M, Moore RA, Derry S, McQuay HJ. Single dose oral tenoxicam for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007591.
- Moore RA, Barden J. Systematic review of dexketoprofen in acute and chronic pain. *BMC Clin Pharmacol.* 2008;8:11.

Moore RA, Derry S, McQuay HJ, Wiffen PJ. Single dose oral analgesics for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2011;(9):CD008659.

Moore RA, Derry S, McQuay HJ. Single dose oral aceclofenac for postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007588.

Moore RA, Derry S, McQuay HJ. Single dose oral acetaminophen for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007589.

Moore RA, Derry S, McQuay HJ. Single dose oral dexibuprofen [S(+)-ibuprofen] for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007550.

Moore RA, Derry S, McQuay HJ. Single dose oral fenbufen for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD007547.

Moore RA, Derry S, McQuay HJ. Single dose oral meloxicam for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD007552.

Moore RA, Derry S, McQuay HJ. Single dose oral sulindac for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD007540.

Moore RA, Derry S, Moore M, McQuay HJ. Single dose oral nabumetone for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD007548.

Moore RA, Derry S, Moore M, McQuay HJ. Single dose oral tiaprofenic acid for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(4):CD007542.

Moore RA, Straube S, Paine J, Derry S, McQuay HJ. Minimum efficacy criteria for comparisons between treatments using individual patient meta-analysis of acute pain trials: examples of etoricoxib, paracetamol, ibuprofen, and ibuprofen/paracetamol combinations after third molar extraction. *Pain.* 2011;152(5):982-989.

Moore RA, Straube S, Wiffen PJ, Derry S, McQuay HJ. Pregabalin for acute and chronic pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007076.

Nalamachu S, Pergolizzi JV, Raffa RB, Lakkireddy DR, Taylor R Jr. Drug-drug interaction between NSAIDs and low-dose aspirin: a focus on cardiovascular and GI toxicity. *Expert Opin Drug Saf.* 2014;13(7):903-917.

Ong CK, Seymour RA, Lirk P, Merry AF. Combining paracetamol (acetaminophen) with nonsteroidal anti-inflammatory drugs: a qualitative systematic review of analgesic efficacy for acute postoperative pain. *Anesth Analg.* 2010;110(4):1170-1179.

Ong KS, Seymour RA. Maximizing the safety of nonsteroidal anti-inflammatory drug use for postoperative dental pain: an evidence-based approach. *Anesth Prog.* 2003;50(2):62-74.

Palmer H, Graham G, Williams K, Day R. A risk-benefit assessment of paracetamol (acetaminophen) combined with caffeine. *Pain Med.* 2010;11(6):951-965.

Roy YM, Derry S, Moore RA. Single dose oral lumiracoxib for postoperative pain in adults. *Cochrane Database Syst Rev.* 2010;(7):CD006865.

Roy YM, Derry S, Moore RA. Single dose oral lumiracoxib for postoperative pain. *Cochrane Database Syst Rev.* 2007;(4):CD006865.

Seib RK, Paul JE. Preoperative gabapentin for postoperative analgesia: a meta-analysis. *Can J Anaesth.* 2006;53(5):461-469.

Silva RC, Riera R, Saconato H. Lumiracoxib for acute postoperative dental pain: a systematic review of randomized clinical trials. *Sao Paulo Med J.* 2011;129(5):335-345.

Skoglund LA. Good evidence for effectiveness of analgesics most commonly prescribed by UK dentists. What is the relative efficacy of single-dose oral analgesics after third molar extraction? *Evid Based Dent.* 2005;6(3):66.

Smith LA, Carroll D, Edwards JE, Moore RA, McQuay HJ. Single-dose ketorolac and pethidine in acute postoperative pain: systematic review with meta-analysis. *Br J Anaesth.* 2000;84(1):48-58.

Steiner TJ, Voelker M. Gastrointestinal tolerability of aspirin and the choice of over-the-counter analgesia for short-lasting acute pain. *J Clin Pharm Ther.* 2009;34(2):177-186.

Straube S, Derry S, Moore RA, Wiffen PJ, McQuay HJ. Single dose oral gabapentin for established acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2010;(5):CD008183.

Sultan A, McQuay HJ, Moore RA, Derry S. Single dose oral flurbiprofen for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007358.

Tirunagari SK, Derry S, Moore RA, McQuay HJ. Single dose oral etodolac for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(3):CD007357.

Toms L, Derry S, Moore RA, McQuay HJ. Single dose oral paracetamol (acetaminophen) with codeine for postoperative pain in adults. *Cochrane Database Syst Rev.* 2009;(1):CD001547.

Toms L, McQuay HJ, Derry S, Moore RA. Single dose oral paracetamol (acetaminophen) for postoperative pain in adults. *Cochrane Database Syst Rev.* 2008;(4):CD004602.

Traa MX, Derry S, Moore RA. Single dose oral fenpropfen for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2011;(2):CD007556.

Wasey JO, Derry S, Moore RA, McQuay HJ. Single dose oral diflunisal for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2010;(4):CD007440.

Wei W, Zhao T, Li Y. Efficacy and safety of parecoxib sodium for acute postoperative pain: a meta-analysis. *Exp Ther Med.* 2013;6(2):525-531.

Weil K, Hooper L, Afzal Z, et al. Paracetamol for pain relief after surgical removal of lower wisdom teeth. *Cochrane Database Syst Rev.* 2007;(3):CD004487.

Wiffen PJ, Derry S, Moore RA, McQuay HJ. Carbamazepine for acute and chronic pain in adults. *Cochrane Database Syst Rev.* 2011;(1):CD005451.

Zemmel MH. The role of COX-2 inhibitors in the perioperative setting: efficacy and safety—a systematic review. *AANA J.* 2006;74(1):49-60.

Zhang J, Ho KY, Wang Y. Efficacy of pregabalin in acute postoperative pain: a meta-analysis. *Br J Anaesth.* 2011;106(4):454-462.

REASON: NOT RELEVANT QUESTION

Ren ZY, Xu XQ, Bao YP, et al. The impact of genetic variation on sensitivity to opioid analgesics in patients with postoperative pain: a systematic review and meta-analysis. *Pain Physician.* 2015;18(2):131-152.

REASON: ROUTE OF ADMINISTRATION NOT ORAL

Bell RF, Dahl JB, Moore RA, Kalso E. Perioperative ketamine for acute postoperative pain. *Cochrane Database Syst Rev.* 2006;(1):CD004603.

Bell RF, Dahl JB, Moore RA, Kalso E. Peri-operative ketamine for acute post-operative pain: a quantitative and qualitative systematic review (Cochrane review). *Acta Anaesthesiol Scand.* 2005;49(10):1405-1428.

Hamilton TW, Athanassoglou V, Mellon S, et al. Liposomal bupivacaine infiltration at the surgical site for the management of postoperative pain. *Cochrane Database Syst Rev.* 2017;2:CD011419.

REASON: NOT SYSTEMATIC REVIEW

Candiotti KA, Yang Z, Rodriguez Y, et al. The impact of CYP2D6 genetic polymorphisms on postoperative morphine consumption. *Pain Med.* 2009;10(5):799-805.
Hegmann KT, Weiss MS, Bowden K, et al. ACOEM practice guidelines: opioids for treatment of acute, subacute, chronic, and postoperative pain. *J Occup Environ Med.* 2014;56(12):e143-e159.
Practice guidelines for acute pain management in the perioperative setting: an updated report by the American Society of Anesthesiologists Task Force on Acute Pain Management. *Anesthesiology.* 2004;100(6):1573-1581.
Richards D. The Oxford Pain Group League table of analgesic efficacy. *Evid Based Dent.* 2004;5(1):22-23.
Spivakovskiy S, Spivakovskiy Y. Reliable evidence for efficacy of single dose oral analgesics. *Evid Based Dent.* 2016;17(2):60-61.
Wong YJ. Single dose oral analgesics for postoperative pain have few adverse events. *Evid Based Dent.* 2016;17(3):83.

APPENDIX 2

References excluded after full manuscript review

REASON: INCLUDED IN A MORE RECENT OVERVIEW

Derry S, Karlin SM, Moore RA. Single dose oral ibuprofen plus codeine for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2015;(2):CD010107.
Derry S, Wiffen PJ, Moore RA. Single dose oral ibuprofen plus caffeine for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2015;(7):CD011509.
Derry S, Wiffen PJ, Moore RA. Single dose oral diclofenac for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2015;(7):CD004768.

REASON: SUPERSEDED BY A MORE RECENT OVERVIEW

Alexander L, Hall E, Eriksson L, Rohlin M. The combination of non-selective NSAID 400 mg and paracetamol 1000 mg is more effective than each drug alone for treatment of acute pain: a systematic review. *Swed Dent J.* 2014;38(1):1-14.
Au AH, Choi SW, Cheung CW, Leung YY. The efficacy and clinical safety of various analgesic combinations for post-operative pain after third molar surgery: a systematic review and meta-analysis. *PLoS One.* 2015;10(6):e0127611.
Moore PA, Hersh EV. Combining ibuprofen and acetaminophen for acute pain management after third-molar extractions: translating clinical research to dental practice. *JADA.* 2013;144(8):898-908.
Moore RA, Derry S, Straube S, Ireson-Paine J, Wiffen PJ. Faster, higher, stronger? Evidence for formulation and efficacy for ibuprofen in acute pain. *Pain.* 2014;155(1):14-21.

REASON: NOT A SYSTEMATIC REVIEW

American Academy of Pediatrics Committee on Psychosocial Aspects of Child and Family Health, American Pain Society Task Force on Pain in Infants, Children, and Adolescents. The assessment and management of acute pain in infants, children, and adolescents. *Pediatrics.* 2001;108(3):793-797.

REASON: INCLUDED STUDIES OF PREOPERATIVE MEDICATION USE

Eipe N, Penning J, Yazdi F, et al. Perioperative use of pregabalin for acute pain: a systematic review and meta-analysis. *Pain.* 2015;156(7):1284-1300.
Mishriky BM, Waldron NH, Habib AS. Impact of pregabalin on acute and persistent postoperative pain: a systematic review and meta-analysis. *Br J Anaesth.* 2015;114(1):10-31.
Nir RR, Nahman-Averbuch H, Moont R, Sprecher E, Yarnitsky D. Preoperative preemptive drug administration for acute postoperative pain: a systematic review and meta-analysis. *Eur J Pain.* 2016;20(7):1025-1043.

REASON: INTERVENTION UNLIKELY TO HAVE A PHARMACOLOGIC BASIS FOR EFFICACY

Raak C, Bussing A, Gassmann G, Boehm K, Ostermann T. A systematic review and meta-analysis on the use of *Hypericum perforatum* (St. John's Wort) for pain conditions in dental practice. *Homeopathy.* 2012;101(4):204-210.

REASON: SYSTEMATIC REVIEWS OF MEDICATIONS NOT MARKETED IN THE UNITED STATES

Bu X, Yang L, Zuo Y. Efficacy and safety of perioperative parecoxib for acute postoperative pain treatment in children: a meta-analysis. *Front Med.* 2015;9(4):496-507.
Derry S, Cooper TE, Phillips T. Single fixed-dose oral dextropropofol plus tramadol for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2016;9:CD012232.
Hearn L, Derry S, Moore RA. Single dose dipyrrone (metamizole) for acute postoperative pain in adults. *Cochrane Database Syst Rev.* 2016;4:CD011421.